

LETTER TO THE EDITOR

Anti-correlation between multiplicity and orbital properties in exoplanetary systems as a possible record of their dynamical histories

A. Zinzi^{1,2} and D. Turrini^{3,4}

¹ Space Science Data Center (SSDC) – ASI, via del Politecnico snc, 00133 Rome, Italy

e-mail: angel.o.zinzi@ssdc.asi.it

² INAF–OAR, via Frascati n. 33, 00078 Monte Porzio Catone (RM), Italy

³ INAF–IAPS, via del Fosso del Cavaliere n. 100, 00133 Rome, Italy

e-mail: diego.turrini@iaps.inaf.it

⁴ Departamento de Física, Universidad de Atacama, 485 Copayapu, Copiapó, Chile

Received 19 July 2017 / Accepted 26 August 2017

ABSTRACT

Context. Previous works focused on exoplanets discovered with the radial velocity (RV) method reported an anti-correlation between the orbital eccentricities of the exoplanets and the multiplicity M (i.e., the number of planets) of the systems they inhabit.

Aims. We further investigate this reported anti-correlation here using a dataset comprising exoplanets discovered with both the RV and transit methods, searching for hints of its causes by exploring the connection between the number of planets and the dynamical state of the exosystems.

Methods. In order to examine the correlation between multiplicity and orbital eccentricity, for every multiplicity case considered ($2 \leq M \leq 6$), we computed the weighted average eccentricities instead of the median eccentricities used previously. The average eccentricities were calculated using the inverse of the uncertainty on the eccentricity values as weights. The analysis of the dynamic state of the exosystems was performed by computing their angular momentum deficit (AMD), which is a diagnostic parameter successfully used in the study of the solar system and recently applied to exosystems as well.

Results. Our results confirm the reported multiplicity-eccentricity anti-correlation and show that the use of the uncertainties on the orbital eccentricities in the analysis allows for a better agreement between the data and the fits. Specifically, our best fit reproduces well the behaviour of the average eccentricities for all systems with $M > 1$, including the additional cases of TRAPPIST-1 ($M = 7$) and of the solar system ($M = 8$). The AMD analysis, while not conclusive due to the limited number of exosystems that could be analysed, also suggests the existence of an anti-correlation between the multiplicity and the AMD of exosystems. This second anti-correlation, if confirmed by future studies, raises the possibility that the population of low-multiplicity exosystems is contaminated by former high-multiplicity systems that became dynamically unstable and lost some of their planets.

Key words. methods: statistical – techniques: radial velocities – planets and satellites: dynamical evolution and stability

1. Introduction

The currently known population of exoplanets shows characteristics that are extremely different from those of their counterparts orbiting the Sun and no exact analogue of the solar system has yet been found (see e.g. [Hatzes 2016](#), for a recent review). Albeit it is likely that our picture of exoplanetary systems could be incomplete because of the limitations of our present observational capabilities, this nevertheless reveals the large variety of exosystems across the Galaxy and the diversity of the outcomes of the processes shaping the formation and evolution of planetary systems (see [Hatzes 2016](#), and references therein).

Exploiting the possibility to perform statistical studies for the first time thanks to the large number of discovered exoplanets, recent works ([Juric & Tremaine 2008](#); [Limbach & Turner 2015](#)) reported the existence of an anti-correlation between the multiplicity of exosystems (i.e. the number of planets they possess) and the orbital eccentricities of their planets. In particular, [Limbach & Turner \(2015\)](#) analysed a sample of 403 exoplanets showing that, once divided in bins according to the multiplicity of their host systems, the medians of their eccentricities can be

fitted by a power law. These authors also verified that the same power law can be used to fit the median of the eccentricities of solar system planets.

In building their sample, [Limbach & Turner \(2015\)](#) selected exoplanets exclusively discovered by means of the radial velocity (RV) method, requiring eccentricity e measurements that are different from zero for all of the planets. This selection criterion ended up in a dataset comprising 276, 81, 25, 12 and 9 exoplanets for multiplicities M equal to 1, 2, 3, 4 and 5, 6, respectively, the last bin comprising two multiplicity values to increase the statistics. Limbach & Turner performed the analysis without considering the uncertainty on eccentricity measurements and calculated a power law fit for the data with $M > 1$; they obtained the relation $e(M) = 0.584M^{-1.20}$, which they found to reproduce well the median eccentricities for $M > 2$.

In the present work we further investigated the findings of [Limbach & Turner \(2015\)](#) by taking into account the uncertainties on the eccentricity measurements in our analysis. To gain a deeper insight on the causes of the observed anti-correlation, we also performed a preliminary analysis of the dynamical state

Table 1. Number of planets in our dataset divided by the multiplicity (M) of their host system.

M	Total	RV	Tr
2	150	140	19
3	44	34	11
4	26	21	12
5	20	14	6
6	18	18	8

Notes. Also reported are the numbers of these planets that have been observed with the RV and transit method.

of the exosystems (instead of the individual exoplanets) in our sample by evaluating their angular momentum deficits (Laskar 2000).

This work is organized as follows: Sect. 2 is dedicated to the description and reduction of the dataset used; in Sect. 3 these data are discussed; and finally, in Sect. 4 conclusions are given.

2. Dataset and methods

To compose our sample we selected planets around stars with effective temperatures between 7920 K and 2600 K (i.e. as expected for type F, G, K, and M stars) in systems with at least two planets. We used the NASA Exoplanet Archive database (Akeson et al. 2013), which we queried by means of the Exoplanetary Analysis and 3D visualization Tool (ExoplAn3T). This web tool, currently under development at the Space Science Data Center (SSDC), allows users to select between the majority of the parameters provided by the NASA Exoplanet Archive application programming interfaces. However, since the query result also includes the required characteristics for all the objects (i.e. star and planets) that constitute the systems, the output of the tool can be used to perform analyses of the systems as a whole.

In our query we made no distinction on the discovery technique of the exoplanets, with the requirement of knowing the estimated uncertainties on their orbital eccentricities. The resulting dataset consisted of 258 exoplanets. Of these, 227 were detected by means of the RV method and 56 were detected with the transit method; the total number is larger than our sample since some exoplanets were observed with multiple techniques. While our sample is smaller than that used by Limbach & Turner (2015), our selection criterion prevents us from introducing further observational biases in the analysis, allowing us to weight the data according to their reliability and information content.

These 258 planets are distributed among the different multiplicities as described in Table 1 and, even though more than half of the planets belongs to systems with $M = 2$, all the considered multiplicity cases possess more than 10 planets, thus providing a statically relevant sample. Our dataset is plotted in Fig. 1, where we also show the planets of TRAPPIST-1 (Gillon et al. 2017) and of the solar system¹. This choice provides us with systems with 7 and 8 planets ($M = 7$ and $M = 8$) respectively, which we used to verify how our results extend to higher multiplicities. These two planetary systems, however, have not been included into our analysis and are shown only for comparison purposes.

2.1. Multiplicity and eccentricity

In analysing the dynamical properties of the exoplanets in our sample, instead of the median values of the eccentricity e in

¹ <https://nssdc.gsfc.nasa.gov/planetary/factsheet/>




Fig. 1. Multiplicity M vs. eccentricity e for all the planets in our sample. The cases for $M = 7$ and $M = 8$ are the TRAPPIST-1 planets and the planets in the solar system. The individual exoplanets in our sample are represented by black crosses. The blue squares represent the mean for every bin for planets observed with the RV method without considering errors. The red triangles are the means associated with planets detected with the RV method taking into account the uncertainties on the eccentricities measurements. Black circles indicate the means for the entire dataset accounting for measurement errors.

the different multiplicity bins considered by Limbach & Turner (2015), we focused on the relation between M and the weighted average eccentricities.

The weights we used in computing the average values of each multiplicity bin were the inverse of the relative errors of the eccentricities, so that

$$\bar{e}(M) = \frac{\sum_{i=1}^M e(M)_i}{\sum_{i=1}^M \frac{1}{\sigma_{e_i}^2}}, \quad (1)$$

where $\bar{e}(M)$ is the weighted average for a specific value of the multiplicity M (from 2 to 6), $e(M)_i$ are the individual eccentricity values of the exoplanets in the multiplicity bin, and σ_{e_i} are the associated relative errors, which in the case of different values for positive and negative errors were computed as

$$\sigma_{e_i} = \frac{\sigma_{e_i}^+ + \sigma_{e_i}^-}{2e_i}. \quad (2)$$

2.2. Multiplicity and dynamical state

Our preliminary investigation of the dynamical state of the exoplanetary systems took advantage of the information supplied by the angular momentum deficit AMD (Laskar 2000), which measures the difference between the angular momentum of a planetary system and the angular momentum the system would have if its planets were all on circular and planar orbits.

As demonstrated by Laskar & Petit (2017), the AMD remains stable after totally inelastic collisions, but increases when a body is ejected from the system. Therefore a system that experienced several ejections (hence decreasing in multiplicity M) should have a higher value of AMD with respect to a system that remained dynamically stable over all its history.

Following Laskar (2000), the AMD is defined as

$$\text{AMD} = \sum_k \frac{p_k}{m_k} \frac{q_k}{Gm_0 a_k} \left(1 - \frac{1}{1 - e_k^2 \cos^2(i_k)} \right), \quad (3)$$

inner system (i.e. the terrestrial planets) showing characteristics of an unstable system and the outer system (i.e. the giant planets) more resembling a stable system.

The analysis of the AMD shows an inverse trend with multiplicity for both the large and the strict selections. We did not attempt to fit the data as the number of systems analysed is statistically limited. Nevertheless, the fact that lower multiplicity systems display a higher value of the AMD, if confirmed when a larger sample of exosystems becomes available, could indicate that high-multiplicity systems are a common outcome of the planetary formation process, but that a significant portion of these systems form or evolves into unstable orbital configurations.

The loss of planets due to phases of dynamical instability by such systems (e.g. by planet-planet scattering events; see Weidenschilling & Marzari 1996; Rasio & Ford 1996) would cause the increase of their AMD values and, consequently, of the eccentricities of their surviving planets. The resulting systems would then contaminate the population of exosystems with lower native multiplicity values, probably explaining the behaviours observed in Figs. 2 and 3.

4. Conclusions

The results of this study confirm the reported anti-correlation between the orbital eccentricity of exoplanets and the multiplicity of the host system they inhabit. The inclusion of the uncertainties on the measured planetary eccentricities in our analysis not only provides a more robust confirmation, but also allows for fitting with a single power law the average eccentricities for multiplicities comprised between $M = 2$ and $M = 6$. Furthermore, both the characteristics of TRAPPIST-1 ($M = 7$) and of the solar system ($M = 8$) are represented well by the same power law

$$e(M) = 0.630M^{-1.02}. \quad (4)$$

In the attempt to provide a physical explanation of the reported behaviour, we made a preliminary analysis of the correlation between the angular momentum deficit AMD of the systems

included in our sample and their multiplicity values. While the data are not sufficient to provide a conclusive answer, we observed an anti-correlation between the multiplicity of the exosystems and their AMD as well. Should this AMD-multiplicity anti-correlation of exosystems be verified once larger and more precise datasets become available, it would provide us with a deeper insight both on the planetary formation process and on the place of the solar system in an exoplanetary context.

Therefore this could be an interesting field of research when the next generation of space telescopes dedicated to exoplanetary systems is ready, starting with the NASA Transiting Exoplanet Survey Satellite (TESS; Ricker et al. 2015), which is planned to be operative in the next year.

Specifically, our results raise the possibility that high-multiplicity systems such as the solar system are a relatively common outcome of the planetary formation process. Our results also suggest that, unlike the solar system, a possibly large portion of these systems form or evolve into dynamically unstable orbital configurations. The ejection of one or more planets from these unstable systems would naturally explain our results by decreasing the multiplicities of the systems increasing their AMD and eccentricities and/or inclinations of their surviving planets at the same time. As a consequence, the population of low-multiplicity exosystems we observe is likely to be contaminated by former high-multiplicity systems that became dynamically unstable and lost some of their planets during their histories.

References

- Akeson, R. L., Chen, X., Ciardi, D., et al. 2013, *PASP*, **125**, 989
 Gillon, M., Triaud, A. H. M. J., Demory, B.-O., et al. 2017, *Nature*, **542**, 456
 Hatzes, A. P. 2016, *Space Sci. Rev.*, **205**, 267
 Juric, M., & Tremaine, S. 2008, in *Extreme Solar Systems*, *ASP Conf. Ser.*, **398**, 295
 Laskar, J. 1997, *A&A*, **317**, L75
 Laskar, J. 2000, *Phys. Rev. Lett.*, **84**, 3240
 Laskar, J., & Petit, A. 2017, *A&A*, **605**, A72
 Limbach, M. A., & Turner, E. L. 2015, *Proc. Nat. Acad. Sci.*, **112**, 20
 Rasio, F. A., & Ford, E. B. 1996, *Science*, **274**, 954
 Ricker, G. R., Winn, J. N., Vanderspek, R., et al. 2015, *J. Astron. Telescopes, Instruments, and Systems*, **1**, 014003
 Weidenschilling, S. J., & Marzari, F., 1996, *Nature* **384**, 619